

## Inertial manifold of the atmospheric equations\*

LI Jianping (李建平)

(State Key Laboratory of Numerical Modelling for Atmospheric Sciences and Geophysical Fluid Dynamics (LASG), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China)

and CHOU Jifan (丑纪范)

(Department of Atmospheric Sciences, Lanzhou University, Lanzhou 730000, China)

Received July 26, 1998

**Abstract** For a class of nonlinear evolution equations, their global attractors are studied and the existence of their inertial manifolds is discussed using the truncated method. Then, on the basis of the properties of operators of the atmospheric equations, it is proved that the operator equation of the atmospheric motion with dissipation and external forcing belongs to the class of nonlinear evolution equations. Therefore, it is known that there exists an inertial manifold of the atmospheric equations if the spectral gap condition for the dissipation operator is satisfied. These results furnish a basis for further studying the dynamical properties of global attractor of the atmospheric equations and for designing better numerical scheme.

**Keywords:** inertial manifold, nonlinear evolution equation, global attractor, operator equation, operator.

Results of qualitative theory on the nonlinear atmospheric dynamics<sup>[1-14]</sup> show that the atmosphere system, whether it is dry or moist, whether there is a topographic effect on it or not, and whether its external forcing is stationary or non-stationary, will evolve to a global attractor as time increases. The long-time behavior of solutions of the system depends on the global attractor, and points out of the attractor, which have nothing to do with the asymptotic state of the system while time tends to infinity, have only transient sense. The existence of the atmosphere attractor reveals that there is a *nonlinear adjustment to external forcing in the system*, and that the asymptotic behavior of solutions of the full atmospheric dynamical equations can be accurately described by a finite-dimensional ordinary differential equation, which furnishes necessary mathematical and physical foundations for setting and designing the models of long-term numerical weather forecast and numerical climate forecast. This attractor, however, may be a fractal and an unsmooth manifold. Moreover, its speed of attraction for orbits of solutions of the system is not exponential. These bring some difficulties for further dynamical analysis and practical computation. Therefore, it is important to find an invariant smooth manifold which contains the global attractor and attracts all the orbits of solutions of the system with exponentially speed. This is another important basic concept of inertial manifold for characterizing asymptotic properties of solutions apart from the concept of global attractor in the studies of infinite-dimensional dynamical system in recent years<sup>[15-17]</sup>. There are significant senses not only for analyzing dynamical properties on global attractor, but also for designing reasonable discretization numerical scheme in

\* Project supported by the Project on Mechanism and Prediction Theory of Heavy Climate Disaster in China and the State Key Project on Dynamics and Prediction Theory of Climate.

investigations on inertial manifold. It is natural to ask whether there exists an inertial manifold in the atmospheric equations or not. The question is mainly discussed in this paper.

## 1 Definition of inertial manifold

*Definition 1.* Assume that the semigroup operator  $\{S(t)\}_{t \geq 0}$  has a global attractor  $A$ . Then subset  $M_I \subset H$  ( $H$  is a Hilbert space) is called an inertial manifold of  $A$  if it satisfies

- (i)  $M_I$  is a finite-dimensional smooth manifold (at least a Lipschitz manifold);
- (ii)  $M_I$  is invariant, i. e.

$$S(t)M_I \subset M_I; \quad (1)$$

(iii)  $M_I$  attracts all the orbits of solutions for semigroup  $\{S(t)\}_{t \geq 0}$  with exponential speed, namely for  $u_0 \in H$  there exist constants  $k_1, k_2 > 0$  such that

$$\text{dist}(S(t)u_0, M_I) \leq k_1 e^{-k_2 t}, \quad \forall t \geq 0; \quad (2)$$

- (iv) the global attractor  $A$  of  $S(t)$  is on  $M_I$ .

## 2 Attractors and manifolds of a class of nonlinear evolution equations

### 2.1 A class of nonlinear evolution equations

Given a Hilbert space  $H$  with inner product  $(\cdot, \cdot)$  and norm  $|\cdot|$ , we study a class of nonlinear evolution equations as follows:

$$\frac{d\vartheta}{dt} + L\vartheta + R(\vartheta) = 0, \quad (3)$$

where

$$R(\vartheta) = \delta_1 B(\vartheta, \vartheta) + \delta_2 B_1(\vartheta, \vartheta) + \delta_3 A\vartheta - f, \quad (4)$$

where  $\delta_i$  ( $i = 1, 2, 3$ ) satisfy

$$\delta_i = 0 \text{ or } 1, \text{ and } \delta_1 + \delta_2 \geq 1. \quad (5)$$

The case for  $\delta_2 = 0$  has been discussed<sup>[15-17]</sup>, but the case for  $\delta_2 = 1$  is first presented and investigated in this paper. The linear operator  $L$  is an unbounded positively definite self-adjoint operator in  $H$ ,  $D(L)$  is dense in  $H$ , and  $L^{-1}$  is compact. The mapping  $\vartheta \rightarrow L\vartheta$  is an isomorphism from  $D(L)$  onto  $H$ . Let  $L^s$  be  $s$  powers ( $s \in \mathbb{R}$ ) of  $L$ , and then the spaces  $V_{2s} = D(L^s)$  are Hilbert spaces with the following inner products:

$$(\vartheta_1, \vartheta_2)_{2s} = (L^s \vartheta_1, L^s \vartheta_2), \quad \forall \vartheta_1, \vartheta_2 \in D(L^s), \quad (6)$$

$\vartheta \in V_s$ . Let

$$|\vartheta|_s = (\vartheta, \vartheta)_s^{1/2}. \quad (7)$$

Since  $L^{-1}$  is a compact self-adjoint operator, it follows from the Rellich inference<sup>[15-17]</sup> that there exist an orthonormal basis  $\{w_j\}_{j=1}^\infty$  and the eigenvalues  $\lambda_j$  of  $L$  in  $H$  such that

$$Lw_j = \lambda_j w_j, \quad (j = 1, 2, \dots) \quad (8)$$

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_j \leq \lambda_{j+1} \leq \dots, \quad (9)$$

$$\lim_{\lambda_j \rightarrow \infty} \lambda_j = +\infty. \quad (10)$$

From (8) and (9), it is easy to obtain

$$|L^{1/2}\vartheta| \geq \lambda_1^{1/2} |\vartheta|, \quad \forall \vartheta \in D(L^{1/2}), \quad (11)$$

$$|L^{s+1/2}\vartheta| \geq \lambda_1^{1/2} |L^s \vartheta|, \quad \forall \vartheta \in D(L^{s+1/2}), \quad \forall s. \quad (12)$$

Letting  $P = P_N$  be an orthogonal projection from  $H$  to  $\text{Span}\{w_1, \dots, w_N\}$ ,  $Q_N = I - P_N$ , one has

$$\lambda_1 \|p\| \leq \|Lp\| \leq \lambda_N \|p\|, \quad p \in PD(L), \quad (13)$$

$$\|Lq\| \geq \lambda_{N+1} \|q\|, \quad q \in QD(L). \quad (14)$$

$B(\vartheta, \vartheta)$  and  $B_1(\vartheta, \vartheta)$  are bilinear operators of  $D(L) \times D(L) \rightarrow H$ , and  $A$  is a linear operator of  $D(L) \rightarrow H$ . Suppose that  $B(\vartheta, \vartheta)$ ,  $B_1(\vartheta, \vartheta)$  satisfy

$$(\delta_1 B(\vartheta, \vartheta_1) + \delta_2 B_1(\vartheta, \vartheta_1), \vartheta_1) = 0, \quad \forall \vartheta, \vartheta_1 \in D(L), \quad (15)$$

$$\|B(\vartheta, \vartheta_1)\| \leq C_1 \|\vartheta\|^{1/2} \|L^{1/2}\vartheta\|^{1/2} \|L^{1/2}\vartheta_1\|^{1/2} \|L\vartheta_1\|^{1/2}, \quad \forall \vartheta, \vartheta_1 \in D(L), \quad (16)$$

$$\|B_1(\vartheta, \vartheta_1)\| \leq C_2 \|L^{1/2}\vartheta\|^{1/2} \|L^{1/2}\vartheta_1\|^{1/2}, \quad \forall \vartheta, \vartheta_1 \in D(L). \quad (17)$$

Then  $A$  satisfies one of the following properties:

$$\|A\vartheta\| \leq C_3 \|L^{1/2}\vartheta\|, \quad \forall \vartheta \in D(L), \quad (18)$$

$$\|A\vartheta\| \leq C_4 \|L^{1/2}\vartheta\|^{1/2} \|L\vartheta\|^{1/2}, \quad \forall \vartheta \in D(L). \quad (19)$$

Additionally,  $B, B_1$  and  $A$  also have the continuity properties:

$$\|L^{1/2}B(\vartheta, \vartheta_1)\| \leq C_5 \|L\vartheta\| \|L\vartheta_1\|, \quad \forall \vartheta, \vartheta_1 \in D(L), \quad (20)$$

$$\|L^{1/2}B_1(\vartheta, \vartheta_1)\| \leq C_6 \|L\vartheta\|^{1/2} \|L\vartheta_1\|^{1/2}, \quad \forall \vartheta, \vartheta_1 \in D(L), \quad (21)$$

$$\|L^{1/2}A\vartheta\| \leq C_7 \|L\vartheta\|, \quad \forall \vartheta \in D(L), \quad (22)$$

where  $C_i$  ( $i = 1, \dots, 7$ ) are positive constants. Besides, assume that  $A + L$  is positively definite, i. e. there exists a constant  $C_8 > 0$  such that

$$((A + L)\vartheta, \vartheta) \geq C_8 \|L^{1/2}\vartheta\|^2, \quad \forall \vartheta \in D(L). \quad (23)$$

If  $A$  is anti-adjoint, i. e.

$$(A\vartheta, \vartheta) = 0, \quad \forall \vartheta \in D(L), \quad (24)$$

then it is obvious that (23) is true. However,  $A$  in (23) may not be an anti-adjoint operator.

## 2.2 Global attractor

We now consider the initial-value problem of (3), namely (3) has the following initial condition:

$$\vartheta(0) = \vartheta_0 \in H. \quad (25)$$

Assume that there exist a unique solution  $S(t)\vartheta_0 \in D(L)$  ( $\forall t \in \mathbb{R}^+$ ) for the initial-value problems (3) and (23). The mapping  $S(t)$  has properties of semigroup as usual.

Based on the properties of the operators mentioned above, one has the following lemma.

**Lemma 1.** For any initial value  $\vartheta_0 \in H$ , there are  $\rho_0, \rho_1$  and  $\rho_2$  which depend on  $\lambda_1, \|f\|$  and  $\|L^{1/2}f\|$  such that

$$\limsup_{t \rightarrow +\infty} \|\vartheta(t)\|^2 \leq \rho_0^2, \quad (26)$$

$$\limsup_{t \rightarrow +\infty} \|L^{1/2}\vartheta(t)\|^2 \leq \rho_1^2, \quad (27)$$

$$\limsup_{t \rightarrow +\infty} \|L\vartheta(t)\|^2 \leq \rho_2^2. \quad (28)$$

Lemma 1 implies that any solution of (3) will run respectively into the following balls after some time  $t \geq \tau > 0$ :

$$B_0 = \{\vartheta \in H, \|\vartheta\| \leq 2\rho_0\}; \quad (29)$$

$$B_1 = \{\vartheta \in D(L^{1/2}), \|L^{1/2}\vartheta\| \leq 2\rho_1\}; \quad (30)$$

$$B_2 = \{\vartheta \in D(L), \|L\vartheta\| \leq 2\rho_2\}. \quad (31)$$

Thus we have Theorem 1.

**Theorem 1.** *There exists a global attractor  $\mathcal{A}$  in (3). It is a limiting set of  $B_2$ , i. e.*

$$\mathcal{A} = \omega(B_2) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} S(t)B_2}, \quad (32)$$

where the closure is taken in  $H$ , and we have

$$\mathcal{A} \subseteq B_2 \cap B_1 \cap B_0. \quad (33)$$

### 2.3 Inertial manifold

Here we discuss inertial manifolds of (3) by use of the truncated method of ref. [15]. Assume that  $\theta(s)$  is a smooth function of  $\mathbb{R}^+ \rightarrow [0, 1]$ :

$$\begin{cases} \theta(s) = \begin{cases} 1, & s \in [0, 1]; \\ 0, & s \geq 2; \end{cases} \\ |\theta'(s)| \leq 2, & s \geq 0. \end{cases} \quad (34)$$

Fix  $\rho = 2\rho_2$ , and define

$$\theta_\rho(s) = \theta(s/\rho), \quad s \geq 0. \quad (35)$$

Then the truncated equation of (3) is

$$\frac{d\vartheta}{dt} + L\vartheta + F(\vartheta) = 0, \quad (36)$$

where  $F(u) = \theta_\rho(|L\vartheta|)R(\vartheta)$ . Clearly, when  $|L\vartheta| \leq \rho$ ,  $\theta_\rho(|L\vartheta|) = 1$ , and in this case (36) is accordant with (3). When  $|L\vartheta| \geq 2\rho$ ,  $\theta_\rho(|L\vartheta|) = 0$ , and in this case making the inner product for the two sides of (36) with  $L^2\vartheta$ , one obtains

$$\frac{1}{2} \frac{d}{dt} |L\vartheta|^2 + \lambda_1 |L\vartheta|^2 \leq \frac{1}{2} \frac{d}{dt} |L\vartheta|^2 + \lambda_1 |L^{3/2}\vartheta|^2 \leq 0. \quad (37)$$

Therefore, orbit  $\vartheta(t)$  is exponentially convergent to the ball whose radius  $\rho_3 \geq 2\rho$  in  $D(L)$ .

Let  $p(t) = P\vartheta$ ,  $q(t) = Q\vartheta$ . Then  $p$  and  $q$  in  $PH$  and  $QH$  satisfy

$$\frac{dp}{dt} + Lp + PF(\vartheta) = 0, \quad (38)$$

$$\frac{dq}{dt} + Lq + QF(\vartheta) = 0, \quad (39)$$

where  $\vartheta = p + q$ , and the inertial manifold  $M_I = \text{Graph}(\Phi)$ , i. e. it is got by the graph of the Lipschitz mapping  $\Phi: PD(L) \rightarrow QD(L)$ . The mapping  $\Phi$  is obtained by the fixed point of the function space  $H_{b,l}$ . The definition of the function space  $H_{b,l}$  is as follows.

**Definition 2.**  $H_{b,l}$  is a function space of the Lipschitz mapping  $\Phi: PD(L) \rightarrow QD(L)$  satisfying

$$|L\Phi(p)| \leq b, \quad b > 0 \text{ is undetermined}, \quad p \in PD(L); \quad (40)$$

$$|L\Phi(p_1) - L\Phi(p_2)| \leq l |Lp_1 - Lp_2|, \quad l > 0, \quad p_1, p_2 \in PD(L); \quad (41)$$

$$\text{supp } \Phi \subseteq \{p \in PD(L) \mid |Lp| \leq 4\rho\}. \quad (42)$$

Introduce

$$\|\Phi_1 - \Phi_2\| = \sup_{p \in D(l)} |L\Phi_1(p) - L\Phi_2(p)|. \quad (43)$$

Then  $H_{b,l}$  is a complete metric space.

For  $\Phi \in H_{b,l}$ , we define an inertial mapping  $T$  of  $PD(L)$ :

$$T\Phi(p_0) = - \int_{-\infty}^0 e^{\tau L} QF(\vartheta) d\tau, \quad p_0 \in PD(L), \quad (44)$$

where  $\vartheta(\tau) = p(\tau; \Phi, p_0) + \Phi(p(\tau; \Phi, p_0))$ ,  $p(\tau; \Phi, p_0)$  is a solution of (36) satisfying  $p(\tau; \Phi, p_0) = p_0$ . Hence, for (3) one has Theorem 2<sup>[12]</sup>.

**Theorem 2.** Suppose that operators  $A, B, B_1$  and  $L$  satisfy (8)—(16), (20)—(23), and (18) (or (19)),  $0 < l < 1/8$ , and that there exist constants  $N_0, K_1$  and  $K_2$  (which depend on  $l$  and initial values) satisfying  $N \geq N_0, \lambda_{N+1} \geq K_1$  and the spectral gap condition  $\lambda_{N+1} - \lambda_N \geq K_2$ . Then there exists  $b > 0$  such that

- (i)  $T$  maps  $H_{b,l}$  into  $H_{b,l}$ ;
- (ii)  $T$  has a fixed point in  $H_{b,l}$ ;
- (iii)  $M_l = \text{Graph}(\Phi)$  is an inertial manifold of (3);
- (iv)  $M_l$  contains the global attractor of (3).

### 3 Inertial manifold of the atmospheric equations

#### 3.1 Description of the equations

In coordinate  $(\lambda, \theta, p)$ , the atmospheric equations of large-scale motion are as follows<sup>[1-3]</sup>:

$$\frac{\partial u}{\partial t} + \Lambda u + \left(2\Omega \cos\theta + \frac{\text{ctg}\theta}{a}u\right)v + \frac{1}{a \sin\theta} \frac{\partial \phi}{\partial \lambda} + L_1 u = 0, \quad (45)$$

$$\frac{\partial v}{\partial t} + \Lambda v - \left(2\Omega \cos\theta + \frac{\text{ctg}\theta}{a}u\right)u + \frac{1}{a} \frac{\partial \phi}{\partial \theta} + L_1 v = 0, \quad (46)$$

$$\frac{\partial \phi}{\partial p} + \frac{R}{p}T = 0, \quad (47)$$

$$\frac{1}{a \sin\theta} \left( \frac{\partial u}{\partial \lambda} + \frac{\partial v \sin\theta}{\partial \theta} \right) + \frac{\partial \omega}{\partial p} = 0, \quad (48)$$

$$\frac{R^2}{C^2} \frac{\partial T}{\partial t} + \frac{R^2}{C^2} \Lambda T - \frac{R}{p} \omega + L_2 T = \frac{R^2}{C^2} \frac{\varepsilon}{C_p}, \quad (49)$$

where

$$\Lambda = \frac{u}{a \sin\theta} \frac{\partial}{\partial \lambda} + \frac{v}{a} \frac{\partial}{\partial \theta} + \omega \frac{\partial}{\partial p},$$

$$L_i = -\frac{\partial}{\partial p} l_i \frac{\partial}{\partial p} - \mu_i \nabla^2 \quad (i = 1, 2),$$

$$l_i = v_i (gp/R\bar{T})^2 \quad (i = 1, 2),$$

$$\nabla^2 = \frac{1}{a \sin\theta} \frac{\partial}{\partial \theta} \sin\theta \frac{\partial}{\partial \theta} + \frac{1}{a^2 \sin^2\theta} \frac{\partial^2}{\partial \lambda^2},$$

$$C^2 = R^2 \bar{T} (\gamma_d - \gamma) / g,$$

$\bar{T} = \bar{T}(p)$  is the mean temperature over isobaric  $p$  surface,  $T$  the deviation from  $\bar{T}$ ,  $\phi$  the deviation from  $\bar{\phi}$ ,  $\varepsilon$  the diabatic heating, and other symbols are meteorologically as usual. The domain of solutions of the equations is  $\Omega = S^2 \times (p_0, P_s)$ ,  $0 < p_0 < P_s < \infty$ , where  $p_0$  is a certain small positive number, and  $P_s$  the atmospheric pressure on the earth's surface. The boundary value conditions are

$$p = p_0, (\partial u / \partial p, \partial v / \partial p, \omega, \partial T / \partial p) = 0, \quad (50)$$

$$p = P_s, u = v = \omega = 0, \quad (51)$$

$$\partial T / \partial p = \alpha_s (T_s - T), \quad (52)$$

where  $T_s$  is the temperature on the earth's surface,  $\alpha_s$  a parameter related to the turbulent thermal conductivity. We only discuss the case of the homogeneous boundary value condition of (52), i.e.

$$\partial T / \partial p = -\alpha_s T. \quad (53)$$

For the non-homogeneous case, it may be discussed by using homogeneous transformation. Here we do not consider the topographic effect, so  $\phi(\lambda, \theta, P_s) = 0$ .

The initial value condition is

$$(u, v, T) |_{t=0} = (u^{(0)}, v^{(0)}, T^{(0)}). \quad (54)$$

From (48) and the boundary value conditions, one has

$$\omega(\lambda, \theta, p) = - \int_p^{P_s} (\partial \omega / \partial p) dp = - \int_p^{P_s} ((\partial u / \partial \lambda + \partial v \sin \theta / \partial \theta) / a \sin \theta) dp. \quad (55)$$

Similarly, one has

$$\phi(\lambda, \theta, p) = - \int_p^{P_s} (\partial \phi / \partial p) dp = - \int_p^{P_s} (RT/p) dp. \quad (56)$$

Equations (45)—(49) are essentially the equations of three variable  $u, v$  and  $T$ , i. e. (45), (46) and (49), where  $\omega$  and  $\phi$  are given by (55) and (56). Therefore, by introducing the vector function

$$\vartheta = (u, v, (R/C)T)^T, \quad (57)$$

eqs. (45)—(49) can be written as the following operator equation:

$$\frac{\partial \vartheta}{\partial t} + R(\vartheta) + L\vartheta = 0, \quad (58)$$

where

$$R(\vartheta) = B(\vartheta)\vartheta + B_1(\vartheta)\vartheta + A\vartheta - \zeta, \quad (59)$$

$$L = \text{diag}(L_1, L_1, C^2 L_2 / R^2), \quad (60)$$

$$B(\vartheta) = \begin{bmatrix} \Lambda_1 & u \text{ctg} \theta / a & 0 \\ -u \text{ctg} \theta / a & \Lambda_1 & 0 \\ 0 & 0 & \Lambda_1 \end{bmatrix}, \quad (61)$$

$$B_1(\vartheta) = \text{diag}\left(\omega \frac{\partial}{\partial p}, \omega \frac{\partial}{\partial p}, \omega \frac{\partial}{\partial p}\right). \quad (62)$$

When  $A$  is a linear operator,

$$A\vartheta = \left( fv + \frac{1}{a \sin \theta} \frac{\partial \phi}{\partial \lambda}, -fu + \frac{1}{a} \frac{\partial \phi}{\partial \theta}, -\frac{C}{p} \omega \right)^T, \quad (63)$$

$$(A\vartheta_1, \vartheta_2) = \int_{\Omega} \left[ \left( fv_1 + \frac{1}{a \sin \theta} \frac{\partial \phi_1}{\partial \lambda} \right) u_2 + \left( -fu_1 + \frac{1}{a} \frac{\partial \phi_1}{\partial \theta} \right) v_2 + \left( -\frac{R}{p} \omega_1 \right) T_2 \right] d\Omega, \quad (64)$$

where  $\omega$  and  $\phi$  are given by (55) and (56), i. e.

$$\omega_i(\lambda, \theta, p) = - \int_p^{P_s} ((\partial u_i / \partial \lambda + \partial v_i \sin \theta / \partial \theta) / a \sin \theta) dp, \quad (i = 1, 2),$$

$$\phi_i(\lambda, \theta, p) = - \int_p^{P_s} (RT_i/p) dp, \quad (i = 1, 2),$$

$$\Lambda_1 = \frac{u}{a \sin \theta} \frac{\partial}{\partial \lambda} + \frac{v}{a} \frac{\partial}{\partial \theta}, \quad (65)$$

$$\xi = (0, 0, R\varepsilon / (CC_p))^T. \quad (66)$$

The diabatic heating  $\varepsilon$  is given and stationary in this paper, and at this same time there exists a unique solution for the initial-boundary value problems (45)—(54)<sup>[5]</sup>. A semigroup  $S(t)$  is therefore defined by the operator equation (58) for the initial value  $\vartheta(0) = \vartheta_0$ .

### 3.2 Properties of operators

Let  $H$  be a Hilbert space (taking  $H = L_2(\Omega)$  in the following parts),  $D(L)$  the domain of definition of operator  $L$  corresponding to the boundary value conditions (50), (51) and (53) (being marked as  $D_{\partial\Omega}$ ). Then  $D(L) = \{\vartheta \mid \vartheta \in C^\infty(\bar{\Omega}), D_{\partial\Omega}\}$ , where  $D(L)$  is a linear dense set.

**Lemma 2.** *Operator  $L$  is symmetrical, namely,*

$$(L\vartheta_1, \vartheta_2) = (\vartheta_1, L\vartheta_2), \quad \forall \vartheta_1, \vartheta_2 \in D(L). \quad (67)$$

**Lemma 3.** *Operator  $L$  is self-adjoint.*

As is easily seen, one has

$$(L\vartheta, \vartheta) \geq 0, \quad (68)$$

for  $\vartheta \in D(L)$ . Furthermore, there exist constants  $C_1, C_2 > 0$  such that

$$(L\vartheta, \vartheta) \geq C_1 \|\vartheta\|_1, \quad (69)$$

$$(L\vartheta, \vartheta) \geq C_2 \|\vartheta\|_0, \quad (70)$$

where

$$\|\vartheta\|_1 = (\|u\|_{H^1}^2 + \|v\|_{H^1}^2 + \|T\|_{H^1}^2)^{1/2}, \quad (71)$$

$$\|\vartheta\|_0 = (\|u\|^2 + \|v\|^2 + \|T\|^2)^{1/2}, \quad (72)$$

where  $\|\cdot\|_{H^1}$  takes the  $H^1(\Omega)$ -norm,  $\|\cdot\|$  the  $L_2(\Omega)$ -norm. Therefore, we have Lemma 4.

**Lemma 4.** *Operator  $L$  is positively definite, i. e. there exists  $C > 0$  such that*

$$(L\vartheta, \vartheta) \geq C(\vartheta, \vartheta), \quad \forall \vartheta \in D(L), \quad (73)$$

where the equality is true if and only if  $\vartheta = 0$ .

Based on the above analyses,  $L$  is a self-adjoint positively definite linear operator. Thus one can introduce a new inner product in  $D(L)$  as follows:

$$(\vartheta_1, \vartheta_2) = (L\vartheta_1, \vartheta_2). \quad (74)$$

This yields a new norm

$$\|\vartheta\|_1 = (\vartheta, \vartheta)_1^{1/2} = (L\vartheta, \vartheta)^{1/2}, \quad \vartheta \in D(L). \quad (75)$$

A new complete Hilbert space can be got by taking the closure of  $D(L)$  according to (75). Obviously,  $D(L)$  is dense in  $H_1$ . Using the positively definite property of  $L$ , one has

$$\|\vartheta\|_0 \leq C^{-1} \|\vartheta\|_1, \quad \vartheta \in D(L). \quad (76)$$

$H_1$  is therefore embedded in  $H$ .

More generally, let  $L^s$  be  $s$  powers ( $s \in \mathbb{R}$ ) of  $L$ , and the spaces  $H_{2s} = D(L^s)$  be the Hilbert spaces with the following inner product:

$$(\vartheta_1, \vartheta_2)_{2s} = (L^s\vartheta_1, L^s\vartheta_2), \quad \forall \vartheta_1, \vartheta_2 \in D(L^s), \quad (77)$$

$\vartheta \in H_s$ . Besides,

$$\|\vartheta\|_s = (\vartheta, \vartheta)_s^{1/2}. \quad (78)$$

**Lemma 5.** *The inverse operator  $L^{-1}$  of operator  $L$  is compact.*

*Proof.* For  $\forall f \in H$ , there exists a unique solution  $\vartheta = L^{-1}f$  of  $L\vartheta = f$  in  $D(L)$ . And using

$$\|\vartheta\|^2 \leq C(L\vartheta, \vartheta) = C(f, \vartheta),$$

one has

$$\|\vartheta\| \leq C^* \|f\|. \quad (79)$$

$L^{-1}$  is therefore a bounded linear operator of  $H \rightarrow H_1$ . In addition, the embedding from  $H_1$  to  $H$  is

compact, so  $L^{-1}$  is compact.

$L^{-1}$  is a self-adjoint compact operator because both  $L$  and  $L^{-1}$  are self-adjoint. It follows that there exists an orthonormal basis  $\{W_j\}_{j=1}^{\infty}$  and the eigenvalues  $\hat{\lambda}_j > 0$  of  $L$  and  $\hat{\lambda}_j > \hat{\lambda}_{j+1}$  such that

$$L^{-1}W_j = \hat{\lambda}_j^{-1}W_j \quad (j = 1, 2, \dots). \quad (80)$$

Let  $\lambda_k = \hat{\lambda}_k^{-1}$ . Then one has

$$LW_j = \lambda_j W_j, \quad W_j \in D(L), \quad (j = 1, 2, \dots), \quad (81)$$

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_j \leq \lambda_{j+1} \leq \dots, \quad (82)$$

$$\lim_{j \rightarrow \infty} \lambda_j = \infty. \quad (83)$$

Therefore,  $L$  has properties the same as those of (11)—(14).

For operators  $B, B_1$  and  $A$  we have Lemma 6.

**Lemma 6.**

$$(B(\vartheta, \vartheta_1) + B_1(\vartheta, \vartheta_1), \vartheta_1) = 0, \quad (84)$$

$$(A\vartheta, \vartheta) = 0. \quad (85)$$

Let

$$B(\vartheta, \vartheta_1) = B(\vartheta)\vartheta_1, \quad (86)$$

$$B_1(\vartheta, \vartheta_1) = B_1(\vartheta)\vartheta_1. \quad (87)$$

Then both  $B(\vartheta, \vartheta_1)$  and  $B_1(\vartheta, \vartheta_1)$  are linear operators of  $D(L) \times D(L) \rightarrow H$ . Thanks to the Holder inequality and the Sobolev embedded inequality, we have Lemma 7.

**Lemma 7.** *There exists a constant  $C_1 > 0$  such that*

$$|B(\vartheta, \vartheta_1)| \leq C_1 \|\vartheta\|_0^{1/2} \|\vartheta\|_1^{1/2} \|\vartheta_1\|_1^{1/2} \|\vartheta_1\|_2^{1/2}, \quad (88)$$

where

$$\|\vartheta_1\|_2 = (\|u\|_2^2 + \|v\|_2^2 + \|T\|_2^2)^{1/2}. \quad (89)$$

According to the Friedrichs inequality, one has

$$\|\omega\| \leq K_1 \|\partial\omega/\partial p\|, \quad (90)$$

where constant  $K_1 > 0$ . Moreover, using the Minkowski inequality, one has

$$\|\omega\| \leq K_1 (\|u\|_1 + \|v\|_1). \quad (91)$$

Similarly, one can get

$$\|\phi\| \leq K_2 \|\partial\phi/\partial p\|, \quad (92)$$

$$\|\phi\| \leq K_2 \|RT/C\|, \quad (93)$$

where constant  $K_2 > 0$ . Additionally, thanks to the Schwarz inequality, we have Lemma 8.

**Lemma 8.** *There exist constants  $k, K > 0$  such that*

$$|B_1(\vartheta, \vartheta_1)| \leq k \|\vartheta\|_1 \|\vartheta_1\|_1, \quad (94)$$

$$|A\vartheta| \leq K \|\vartheta\|_1. \quad (95)$$

### 3.3 Inertial manifold

It is easy to see that Lemma 1 is true for eq. (58) because operators in (58) satisfy the properties of operators in (3) according to lemmas 2—8. Furthermore, Theorem 2 is true for eq. (58) if the spectral gap condition is satisfied. This shows that there exist an global attractor and an inertial manifold  $M_I$  in the atmospheric equations (45)—(49) under that condition, and the inertial manifold contains the attractor and attracts all the orbits of solutions of eqs. (45)—(49) with exponential speed.



## 4 Conclusion

Global attractors and inertial manifolds of a class of nonlinear evolution equations are studied in this paper. And it is proved that the atmospheric equations just belong to this kind of nonlinear evolution equations, and that there exists an inertial manifold of the atmospheric equations if the spectral gap condition for the dissipation operator is satisfied. These results furnish a basis for understanding further the dynamical construction and characteristic of the global attractor of the atmosphere because the inertial manifold is an invariant smooth manifold which attracts all the orbits of solutions with exponential speed. Additionally, we may use the existence of inertial manifold to design a better numerical scheme to simulate precisely the long-term variation of the atmosphere. If there exists an inertial manifold  $M_{\tau_I}$  (here  $\tau$  is the time stepsize) in the discretization numerical scheme designed and  $M_{\tau_I} \rightarrow M_I$  as  $\tau \rightarrow 0$ , this kind of numerical scheme is able to simulate precisely the long-term behavior of solutions of the original equations. This will be reported in another paper.

## References

- 1 Chou Jifan, Some properties of operators and the decay of effect of initial condition, *Acta Meteor. Sinica* (in Chinese), 1983, 41(4): 385.
- 2 Chou Jifan, *Long-term Numerical Weather Prediction* (in Chinese), Beijing: China Meteorological Press, 1986, 51—95.
- 3 Chou Jifan, Some general properties of the atmospheric model in  $H$  space,  $R$  space, point mapping, cell mapping, in *Proceedings of International Summer Colloquium on Nonlinear Dynamics of the Atmosphere*, 10—20 Aug, Beijing: Science Press, 1986, 187-189.
- 4 Wang Shouhong, Huang Jianping, Chou Jifan, Some properties of solutions for the equations of large-scale atmosphere, nonlinear adjustment to the time-independent external forcing, *Science in China*, Ser. D, 1990, 33(4): 476.
- 5 Chou Jifan, Global analysis of the dynamical equations of atmospheric motion, *J. Beijing Meteor. College* (in Chinese), 1995, (1): 1.
- 6 Chou Jifan, Some advances and trends in the atmospheric dynamics, *The Forward Position and Prospects for Modern Atmospheric Sciences* (in Chinese), Beijing: China Meteorological Press, 1995, 71—75.
- 7 Li Jianping, Chou Jifan, The property of solutions for the equations of large-scale atmosphere with the non-stationary external forcing, *Chinese Science Bulletin*, 1996, 41(7): 587.
- 8 Li Jianping, Chou Jifan, Existence of atmosphere attractor, *Science in China*, Ser. D, 1997, 40(2): 215.
- 9 Li Jianping, Chou Jifan, Effects of external forcing, dissipation and nonlinearity on the solutions of atmospheric equations, *Acta Meteor Sinica* (in Chinese), 1997, 11(2): 57.
- 10 Li Jianping, Chou Jifan, Further study on the properties of operators of atmospheric equations and the existence of attractor, *Acta Meteor Sinica* (in Chinese), 1997, 11(2): 216.
- 11 Li Jianping, Qualitative theory of the dynamical equations of atmospheric and oceanic motion and its applications, *Ph. D Dissertation* (in Chinese), Lanzhou: Lanzhou University, 1997, 209.
- 12 Li Jianping, Chou Jifan, Asymptotic behavior of solutions of the moist atmosphere, *Acta Meteor Sinica* (in Chinese), 1998, 56(2): 187.
- 13 Li Jianping, Chou Jifan, Qualitative theory on the dynamical equations of atmospheric motion and its applications, *Scientia Atmospherica Sinica* (in Chinese), 1998, 21(4): 443.
- 14 Foias, C., Sell, G. R., Temam, R., Inertial manifolds for nonlinear evolutionary equations, *J. Diff. Eqa.*, 1988, 73: 309.
- 15 Constantin, P., Foias, C., Temam, R., *Attractors Representing Turbulent Flows*, USA: Memoirs Amer. Math. Soc., 1985, 314.
- 16 Guo Buolin, *Nonlinear Evolutionary Equations* (in Chinese), Shanghai: Shanghai Scientific and Technological Press, 1995, 183—343.
- 17 Li Kaitai, Ma Yichen, *Hilbert Space Methods for Equations of Mathematical Physics* (in Chinese), Xi'an: Xi'an Jiaotong University Press, 1992, 267—400.